

# Effective Utilization of Black Soldier Flies Larvae (*Hermetia illucens*) as a poultry feed: a review

1<sup>st</sup> U.G Binuka Bashura

Department of Agricultural Technology  
SLTC Research University  
Padukka 1 10500, Sri Lanka.  
[binukasenadeera@gmail.com](mailto:binukasenadeera@gmail.com)

2<sup>nd</sup> Sisitha Ramachandra

Department of Agricultural Technology  
SLTC Research University  
Padukka 1 10500, Sri Lanka.  
[\\*sisithar@sltc.ac.lk](mailto:*sisithar@sltc.ac.lk)

3<sup>rd</sup> Amali U. Alahakoon

Department of Biosystems Technology  
University of Sri Jayewardenepura  
Pitipana, Hormagama, Sri Lanka.  
[amalialahakoon@sjp.ac.lk](mailto:amalialahakoon@sjp.ac.lk)

**Abstract** - *The use of food that can be consumed by humans as animal feed has become a major problem in today's world. In 2020, one out of every three people will fail to get enough food on a daily basis. Therefore, the carbohydrates, vitamins, and micronutrients essential for physical activity provided by grains have become unavailable in humans. Consequently, it is crucial for the global community to prioritize the development of substitutes for poultry, and other animal feeds. (BSF) The larval stage of this insect offers a remarkable source of protein, presenting a potential path to replace conventional poultry feed derived from grain resources. Research is being conducted on the use of this protein sources, and it is important to solve these problems and produce animal feed that can be obtained by the animal without shortage and at a low cost to the animal keeper. This offers an optimal remedy for countries such as Sri Lanka, which heavily rely on foreign imports for animal feed. This review emphasizes the introduction of the BSF as a means of developing an equitable resolution, not only for Sri Lanka but also for the global population of chicken and other poultry species.*

**Keywords** - *Black Solider Fly Larvae (BFSL), Animal feed ingredient, Grains, Poultry, Sri Lanka.*

## I. INTRODUCTION

The poultry sector is a fast-growing and flexible industry that is essential for the livelihood of resource-poor farmers. It contributes to improving human nutrition, generating income, and providing manure for crop production. The sector has experienced significant growth and plays a crucial role in the livelihoods of one billion of the world's poorest people [1]. However, in the global economic context, using grains intended for human consumption as animal feed is not a justifiable practice. Approximately one-third of the annual global grain production is designated exclusively for animal feed [2]. The majority of poultry farm owners worldwide continue to use commercial chicken feed without any apparent decrease. In developing countries, large poultry farms are frequently established by feed companies, making them reliant on commercial poultry feed right from the start. This dependence on imported poultry feed or its

components become a noticeable economic challenge in resource-constrained regions like Sri Lanka when trying to secure essential nourishment [3], [4]. Instead of depending on imported feeds, it's more cost-effective to create our own animal feed using affordable ingredients. Protein is generally the most expensive feed ingredient in poultry feed. Contemporary poultry feeds primarily comprise grains, soybean oil meal, minerals, and vitamins as protein supplements [5]. Hence, the BSF emerges as an elegant solution to conserve essential grains for the burgeoning global population while furnishing a protein source for poultry [6]. The imperative establishment of a sector centered on BSF larvae -based feed underscores the growing significance of this innovative approach in the animal feed industry. However, to realize its substantial potential, a conducive environment that entails enhanced government support, cost-effective investment, minimized operational expenses, improved regulatory coordination and heightened consumer acceptance is required [7].

The multifaceted utility of the BSF extends beyond its application as an animal feed. This remarkable insect has already been harnessed for organic waste management on a global scale [8], offering substantial potential as a viable solution for addressing the staggering 1.3 billion metric tons of food waste [9]. This helps minimize the negative impacts of such practices, including air and water pollution, land degradation, methane and hazardous leachate emissions, and climate change [10]. Exploring this topic, the aim of this article is to demonstrate how the use BSF can significantly improve the feed supply in the poultry industry.

## II. BLACK SOILDER FLY

The BSF, scientifically known as *Hermetia illucens* (Linnaeus, 1758), (Diptera: Stratiomyidae), is a synanthropic and polysaprophagous insect indigenous to the Neotropics. Over the years, it has undergone extensive global dispersion, extending its habitat to virtually all zoogeographic regions, particularly in warmer climates [11] [12]. BSF belong to the Stratiomyidae family, which encompasses approximately 2700 species [13]. These flies adults characterized by their medium size and

predominantly black coloration, with certain body parts displaying can range from combinations of metallic reflective colors such as blue-green [14]. Adults have a compact head, notably slenderer than the body, with widely separated eyes. Its head is shiny black, with a white stripe running along the lower facial region banking the eyes. The dark brown to black antennae have eight segments, with the final segment flattening. The BSF has a sponge-like, liquid-feeding mouthpart, which is typical of non-biting flies [15,16,17]. Male and female adults have unexplored transparent 'windows' on their first abdominal segment. The function of which has not yet been definitively studied, but these may serve several purposes, including thermoregulation and communication, or simply be a result of the structure of its exoskeleton. Adults measure 16-17 mm in length, 2.8-3.8 mm in width and 0.220 g (low) in mass [18]. Sexual dimorphism is evident in BI, with females typically larger than males. Females display varying patterns on the head and spot sizes on the abdomen. Male genitalia, the aedeagal complex, are slender with basal dilation. Female genitalia have external features known as terminalia, which are characterised by long cerci segmented into two parts (Scissor-like structure). The genital furca exhibits a subtriangular shape with proximal pointed ends, a significant median aperture, and broad, leaf-shaped posterolateral projections [19,20].

### III. IMPORTANCE OF BSF IN POULTRY FEED

An immediate imperative exists for Sri Lanka and the global community to seek alternative sources of animal feed, distinct from traditional grains. The consequence of not doing so may entail a future marked by food scarcity affecting both human and animal populations. This urgency arises not only from population growth but also from the escalating impact of daily worsening weather and climate changes worldwide.

Until recently, eggs were the cheapest source of animal protein available for human consumption in Sri Lanka. However, since 2022, it has escaped the "cheap" label because the price of an egg has increased to Rs.62 per egg since August 2023, and the price of a kilo of chicken has increased to Rs.1300 [21]. This is a huge amount of money even if you look at the daily expenses of the poor people of Sri Lanka and the common people. The primary factors behind this situation were the insufficient "Maha season" maize harvest in Sri Lanka and the substantial expenses

incurred by chicken feed producers due to maize imports. Consequently, the sharp rise in egg prices, in particular, triggered a significant shift from direct consumption to mass-scale bakery products. This shift, coupled with the subsequent unavailability of the most economical protein source, acted as a catalyst for malnutrition [22].

#### A. *As a high-protein poultry feed.*

BSF larval meal emerges as a promising alternative for nutrient-rich and sustainable animal feed, replacing costly and limited protein sources like fishmeal and soybean meal. The insects considered appropriate for animal feed should be capable of large-scale industrial rearing, producing at least 1,000 kg insect fresh weight daily [23]. When quantifying the crude protein content within BSF dried powder (100 g), it is expressed within a specific range, typically falling between 39% and 49% based on research findings [24]. The variability in this range is attributed to the substrate used. According to the available source data, the median crude protein content was approximately  $45.01 \pm 0.19\%$  [25].

Today, poultry farming mainly focuses on egg-laying hens and broilers for meat production [26]. The protein content in the diet for broiler (Table 1) and egg - laying hens (Table 2) fluctuate based on their age. A young laying hen typically consumes approximately 100–150g [27] of poultry feed and a five-week-old broiler consumes approximately 1200–1300 g [28]. According to the table data, only approximately 50% of BFS is required to provide the daily protein requirement of chickens.

TABLE I.

Protein requirement [29] of broilers (Ross 308) according to their age and the protein percentage of chicken feed for the respective age in the world [30] and in the Sri Lankan [31] market (100g).	Age (days)		
	1 - 10	11 - 24	25 - slaughter
Protein requirement of broiler per day (min %)	22 – 25	21 – 23	19 – 21
Protein content of traditional broiler chicken feed in the global market (min %)	21.95	20.92	19.90
Protein content of broiler chicken feed in the Sri Lanka (min %)	23.0	21.5	19.5

TABLE II.

Protein requirement [32] EU red laying hens, according to their age and the protein percentage of poultry feed [33] for the respective age in the global market (100g).	Age (weeks)			
	1 - 5	6 - 10	11 -16/17	18 - 50
Protein requirement of red egg laying breed per day (min %)	20	20	16	18
Protein content of traditional layer chicken feed in the global market (min %)	19.09	17.07	15.38	18.25

### B. Supplementing other nutritional needs of poultry.

Apart from proteins, there are also other nutrient and vitamins that are required for poultry[34,35].They are also mostly contained in BSFL (table 3).

TABLE III.

Daily nutritional requirements of broiler and laying hens and dry BSFL nutritional value.	For broilers between 11 and 24 days old	for laying hens between 18 and 50 weeks old	Dried BSFL powder (100g)
Crude fibre, % (Max.)	5.00	9.00	8.95
Ash, % (Max.)	2.50	4.00	7.91

Calcium, % (Min.)	1.0	3.00	21.00
Phosphorous % (Min.)	0.45	0.50	1.91
Lysine, % (Min.)	1.20	0.70	2.97
Methionine, % (Min.)	0.50	0.35	0.87
Methionine + Cystine %	0.90	0.60	NA
Metabolizable Energy (Kcal/Kg) Min.	3000	2600	2600 – 3500**

Amino acids are crucial for sustaining physiological functions in animals. Some amino acids are categorized as essential because the animal body cannot synthesize them and must acquire them through the diet. These include cysteine, histidine, isoleucine, leucine, lysine, methionine, phenylalanine, threonine, tryptophan, tyrosine, and valine. Non-essential amino acids, such as glutamine, glutamate, proline, glycine

and arginine, can be synthesized by the animal body from other amino acids. Insect proteins are highly sought after for their well-balanced and adaptable amino acid profiles. BSF represent a promising insect protein source for animal feed. Nevertheless, BSF may lack specific essential amino acids (Methionine + Cystine) and may require supplementation with other protein sources or synthetic amino acids to achieve an optimal balance in diet formulations and fulfill amino acid requirements.

BFSL exhibit an approximate lipid content of 30% on a dry matter basis [36], and this lipid composition varies depending on the specific growth media employed. The utilization of BSF in food waste environments led to a prepupae fatty acid composition prominently characterized by a high concentration of lauric acid. This abundance of lauric acid [37] is notably elevated in larvae cultivated on fruit-based substrates (52%) [38]. Larvae demonstrates the capacity to biosynthesize lipids using the available carbohydrates within the substrate [39]. The protein content of younger larvae, harvested at an early developmental stage, is notably elevated, whereas as the larvae matures, their lipid content proportionally increases [40,41]. Fatty acids of BSF larvae grown in a fruit substrate medium were as follows: Lauric Acid (C12:0) – 76.13, Myristic Acid (C14:0) -8.46, Palmitic Acid (C16:0) – 6.98, Oleic Acid (C18:1n-9) – 4.97 [42].

Considering the aforementioned facts, BSF can be recommended as a viable option to supply both macro and micro nutrients while offering it as a preferred feed for poultry. Insect eating is a bird’s natural behavior Birds are omnivorous and like to catch and eat more live things. Feeding live larvae to chickens elicits increased feed intake and weight gain due to the larvae’s motion capturing the birds’ attention [43].

*C. The cost of poultry feed can be greatly reduced*

The global poultry feed market, valued at USD 175.9 billion in 2018 with a 4.5% CAGR, is primarily driven by the increasing demand for chicken meat and eggs worldwide. Poultry is an economical source of protein, and its demand is influenced by population and income levels. Packaged poultry feed is growing faster than traditional feed, offering new opportunities.

Quality and pricing of the feed are significant factors. Key players include Cargill, Archer Daniels, Midland, Kemin Industries, and Lallemand, Inc. In the livestock sector, broilers account for over 65% of the market, primarily for meat production, while layers, contributing about 30%, are reared for egg production. High consumption of eggs and population growth drive the layer segment. Regions such as the United States, Mexico, and Canada are prominent markets for poultry products. The diminishment of arable land, water scarcity, fertilizer constraints, climatic perturbations, and geopolitical conflicts will likely exacerbate global grain prices. Consequently, there will be a commensurate escalation in the cost of animal feed.

Due to the increase in the price of animal feed in the Sri Lankan market, the purchase of products made by turning eggs and meat into ingredients has increased. Today, in the Sri Lankan market, layer starter (25 kg) – Rs.6,071.50, layer grower (25 kg) – Rs.4895, layer special (50 kg) – Rs.9625, broiler booster (50 kg) – Rs.14,615.50, broiler starter (50 kg) – Rs.14,408, broiler finisher (50 kg) – Rs.14,201.50 (figure 01).

PRODUCT CODE	PRODUCT	BAG SIZE	FORM	EX FACTORY PRICE PER BAG WITHOUT TAXES RS.	VAT 15% RS.	EX FACTORY PRICE PER BAG WITH TAXES RS.	EX FACTORY PRICE PER TON WITH TAXES RS.
<b>BROILER FEEDS</b>							
2761	BOOSTER	(50 Kg)	CRUMBLE	12,709.13	1,906.37	14,615.50	292,310.00
2771	STARTER	(50 Kg)	CRUMBLE	12,528.70	1,879.30	14,408.00	288,160.00
2791	FINISHER	(50 Kg)	PELLET	12,349.13	1,852.37	14,201.50	284,030.00
2792	WITHDRAWAL	(50 Kg)	PELLET	12,323.04	1,848.46	14,171.50	283,430.00
<b>LAYER FEEDS</b>							
2831	LAYER STARTER	(25 Kg)	CRUMBLE	5,279.57	791.93	6,071.50	242,860.00
2854	ACCO LAYER GROWER	(25 Kg)	CRUMBLE	4,256.52	638.48	4,895.00	195,800.00
2856	SPECIAL LAYER GROWER	(25 Kg)	CRUMBLE	4,021.74	603.26	4,625.00	185,000.00
2880	ACCO LAYER	(50 Kg)	MASH	8,770.87	1,315.63	10,086.50	201,730.00
2882	ACCO LAYER	(50 Kg)	CRUMBLE	8,628.26	1,294.24	9,922.50	198,450.00
2884	ACCO LAYER SPECIAL	(50 Kg)	CRUMBLE	8,369.57	1,255.43	9,625.00	192,500.00
<b>CATTLE FEEDS</b>							
VAT Exempted							
4010	CALF STARTER	(25 Kg)	PELLET	3,170.00	-	3,170.00	126,800.00
4050	MILK PLUS	(25 Kg)	PELLET	3,350.00	-	3,350.00	134,000.00
4051	MILK STAR	(25 Kg)	PELLET	2,457.63	-	2,457.63	98,305.00
<b>HERBIVORE FEEDS</b>							
4050 ZH	HERBIVORE FEEDS	(25 Kg)	PELLET	4,342.50	-	4,342.50	173,700.00
<b>RABBIT FEEDS</b>							
VAT Exempted							
6001	RABBIT FEED	(50 Kg)	PELLET	8,675.00	-	8,675.00	173,500.00
<b>PIG FEEDS</b>							
VAT Exempted							
5041	PIG FATTENER	(25 Kg)	MASH	5,016.00	-	5,016.00	200,640.00
5050	SOW GESTATION	(25 Kg)	MASH	4,558.25	-	4,558.25	182,330.00
5060	SOW LACTATION	(25 Kg)	MASH	4,037.25	-	4,037.25	161,490.00

Fig. 1. Lives stock feed price list October 2023

However, BSFL can save this amount and grow it on wasted kitchen waste and crop residues, thus managing waste and making poultry farming more profitable. In addition, because this larva can be raised in vertical racks in a small amount of space, the production of one type can be developed as an additional source of income for the poultry farmer. It is easy to use because it is not toxic with other foods, and this fly can be found in more countries around the world. Moreover, it is easy for any common poultry farmer to start this business, there is a high demand for BSFL protein in the world market, it can be used to fill the protein needs of humans [44]. BSF has a short life span and large body size, and has very high feed conversion efficiency [45]. Using BSFL is a highly cost-effective approach that has already been successfully implemented. Positive outcomes from these endeavours offer compelling evidence applicable to both developing and developed nations. Indeed, this is a "**Black diamond**" in the animal feed industry.

#### IV. BSFL GROWING STEPS FOR SRI LANKA

Sri Lanka is, classified as a developing nation, as it heavily relies on importing substantial quantities of grain for plant-based protein as animal feed, as elaborated in the fourth section of this article. The protein content typically falls within the range of 10% to 15%, with the highest concentration of protein [46] residing in the storage proteins, such as prolamins, globulins, and germins [47]. Because of that Poultry farmers need to be informed about the potential establishment of BSF as a viable solution to address Sri Lanka's economic challenges stemming from significant expenditures on grain imports from other nations and the resultant surge in egg prices due to escalating costs of chicken feed. It is imperative to emphasize that relying solely on egg imports from India is not a sustainable resolution [48]. When establishing a BSF rearing facility, whether on a large or small scale, the initial step involves the construction of an adult fly rearing enclosure, often referred to as a "mating cage" or "love cage. On a smaller scale, this enclosure can be crafted from materials like wooden framing with a fine nylon mesh [49] covering, meticulously designed to prevent fly escape. Subsequently, it is crucial to introduce female and male flies into the enclosure in a controlled ratio of 40% females to 60% males. This controlled environment promotes successful mating and egg production in a BSF rearing operation [50]. Adult BSF

flies do not require solid food; they only require water. To enhance their vitality, providing a water source with a slight sugar solution can be beneficial [51]. It is essential to ensure that the water source is provided in a manner that avoids direct contact with the flies, as any physical disturbance may lead to accidental immersion and mortality in the water. The female BSF typically lays an egg cluster containing 540 to 1505 eggs [52], displaying a distinct yellow-cream colouration. When laying these eggs, the BSF seeks a suitable crevice in close proximity to the organic matter. The eggs are meticulously placed within the Substrate and this process can be aided by using a wooden strip or cardboard.

This process involves the construction of a structure capable of accommodating organic material within a tray. The eggs, measuring approximately 1 mm in length, are incubated for approximately 4 days to produce larvae. During the first four days, these larvae exhibit a cream-white colouration and experience their most rapid growth within an approximately 11-day span. Male larvae typically attain a length ranging between 17 and 18 mm, whereas their female counterparts achieve lengths of 21-23 mm. Their diet primarily consists of organic sources such as vegetables, fruits, fish waste, kitchen scraps, fibrous waste, almond hulls, rice straw, high-moisture feed, and livestock and poultry manure [53]. After the 13 to 18 days good for harvest larvae, this phase referred to as the "prepupae" period. During this preparatory phase, the larvae consume an amount of food equal to twice their body size. Subsequently, the prepupa are subjected to a drying process, which can be achieved through mechanical or high-temperature methods to retain moisture content, ultimately converting them into a powder. This powder can be used for both animal and human consumption, and live larvae can be employed as feed for free-ranging poultry. Furthermore, scientific inquiries are underway to examine various proportions and formulations of chicken feed blended with BSFL for improved sustainability and resource efficiency in poultry nutrition [54].

#### V. CONCLUSION

The use of BSFL (*Hermetia illucens*) as poultry feed presents a promising and sustainable approach to enhance global and Sri Lankan grain preservation for human consumption. As the world faces increasingly dire issues of grain wastage for poultry feeds, such innovative solutions are urgently needed to address the problem. By harnessing the potential of BSFL as alternative protein and nutrient sources for poultry, this strategy not only reduces the pressure on traditional grain resources but also aligns with the imperative to minimize grain wastage. Moreover, this eco-friendly approach underscores the significance of interdisciplinary scientific collaboration, offering a potential blueprint for a more resilient and sustainable future in agriculture and food production. Further research in this area is essential to uncover additional nuances and optimize the utilization of

these resources to maximize their impact on grain preservation and food security.

## REFERENCES

- [1] FAO. (2017). The State of Food and Agriculture 2017: Leveraging food systems for inclusive rural transformation. The Food and Agriculture Organization of the United Nations.
- [2] FAO. (2022). The State of World Agriculture 2022: How to make agricultural and food systems more resilient to shocks and crises. The Food and Agriculture Organization of the United Nations.
- [3] Food and Agriculture Organization of the United Nations. (2013). Poultry Development Review. Retrieved from <https://www.fao.org/3/i3531e/i3531e.pdf>
- [4] Food and Agriculture Organization of the United Nations. (2013). Poultry Development Review. Retrieved from <https://www.fao.org/3/i3531e/i3531e.pdf>
- [5] FAO. (2022). The State of World Agriculture 2022: How to make agricultural and food systems more resilient to shocks and crises. Retrieved from <https://www.fao.org/3/cc0451en/cc0451en.pdf>
- [6] FAO. (2022). Insects as food and feed: Opportunities and challenges for sustainability. Retrieved from <https://www.fao.org/3/cb6350en/cb6350en.pdf>
- [7] Food and Agriculture Organization of the United Nations. (2013). Poultry Development Review. Retrieved from <https://www.fao.org/3/i3531e/i3531e.pdf>
- [8] Amrul, N. F., Kabir Ahmad, I., Ahmad Basri, N. E., Suja, F., Abdul Jalil, N. A., & Azman, N. A. (2022). A Review of Organic Waste Treatment Using Black Soldier Fly (*Hermetia illucens*). *Sustainability*, 14(8), 4565. <https://doi.org/10.3390/su14084565>
- [9] The World Counts. (2023). Food Waste Facts. Retrieved from <https://www.theworldcounts.com/stories/food-waste-facts>
- [10] Intergovernmental Panel on Climate Change (IPCC). (2023). Climate Change 2023: Synthesis Report. Retrieved from <https://www.unep.org/resources/report/climate-change-2023-synthesis-report>
- [11] Van Huis, A., & Tomberlin, J. K. (2015). Black soldier fly larvae: biology and utilization for the valorization of organic waste streams. *Insect Science and Its Application*, 36(1), 1-12.
- [12] Amrul, N. F., Kabir Ahmad, I., Ahmad Basri, N. E., Suja, F., Abdul Jalil, N. A., & Azman, N. A. (2022). A Review of Organic Waste Treatment Using Black Soldier Fly (*Hermetia illucens*). *Sustainability*, 14(8), 4565. <https://doi.org/10.3390/su14084565>
- [13] Wikipedia. (2023, October 17). Stratiomyidae. Retrieved from <https://en.wikipedia.org/wiki/Stratiomyidae>
- [14] University of Florida Institute of Food and Agricultural Sciences. (2023, October 17). Black soldier fly: A guide to production and use. Retrieved from <https://edis.ifas.ufl.edu/publication/IN830>
- [15] Kim, W., Bae, S., Park, H., Park, K., Lee, S., Choi, Y., Han, S., & Koh, Y. (2010). The larval age and mouth morphology of the black soldier fly, *Hermetia illucens* (Diptera: Stratiomyidae). *International Journal of Industrial Entomology*, 21, 185-187.
- [16] Oliveira, F., Doelle, K., & Smith, R. (2016). External morphology of *Hermetia illucens* Stratiomyidae: Diptera (L.1758) based on electron microscopy. *Annual Research & Review in Biology*, 9(5), 1-10.
- [17] Rozkosny, R. (1982). A biosystematic study of the European Stratiomyidae (Diptera). Dr W. Junk Publishers, The Hague: Series Entomologica.
- [18] Rozkosny, R. (1982). A biosystematic study of the European Stratiomyidae (Diptera). Dr W. Junk Publishers, The Hague: Series Entomologica.
- [19] Kim, W., Bae, S., Park, H., Park, K., Lee, S., Choi, Y., Han, S., & Koh, Y. (2010). The larval age and mouth morphology of the black soldier fly, *Hermetia illucens* (Diptera: Stratiomyidae). *International Journal of Industrial Entomology*, 21, 185-187.
- [20] Sheppard, D., Tomberlin, J., Joyce, J., Kiser, B., & Sumner, S. (2002). Rearing methods for the black soldier fly (Diptera: Stratiomyidae). *Journal of Medical Entomology*, 39(4), 695-698.
- [21] Tomberlin, J., & Sheppard, D. (2001). Lekking behavior of the black soldier fly (Diptera: Stratiomyidae). *Florida Entomologist*, 84(4), 729-730.
- [22] Economy Next. (2023, October 21). Sri Lanka chicken, egg prices soar due to demand spike, animal feed shortage. Retrieved from <https://economynext.com/sri-lanka-chicken-egg-prices-soar-due-to-demand-spike-animal-feed-shortage-98486/>
- [23] Central Bank of Sri Lanka. (2022). Annual report 2022. Retrieved from [https://www.cbsl.gov.lk/sites/default/files/cbslweb\\_documents/publications/annual\\_report/2022/en/13\\_Box\\_05.pdf](https://www.cbsl.gov.lk/sites/default/files/cbslweb_documents/publications/annual_report/2022/en/13_Box_05.pdf)
- [24] Huis, A. van. (2013). Potential of insects as food and feed in assuring food security. *Annual Review of Entomology*, 58, 563-583. <https://doi.org/10.1146/annurev-ento-120811-153704>
- [25] Zulkifli, N. F., Seok-Kian, A. Y., Seng, L. L., Mustafa, S., Kim, Y. S., & Shapawi, R. (2022). Nutritional value of black soldier fly (*Hermetia illucens*) larvae processed by different methods. *PLOS One*, 17(2), e0263924. doi: 10.1371/journal.pone.0263924
- [26] Zulkifli, N. F., Seok-Kian, A. Y., Seng, L. L., Mustafa, S., Kim, Y. S., & Shapawi, R. (2022). Nutritional value of black soldier fly (*Hermetia illucens*) larvae processed by different methods. *PLOS One*, 17(2), e0263924. doi: 10.1371/journal.pone.0263924
- [27] Food and Agriculture Organization of the United Nations. (2013). Poultry Development Review. Retrieved from <https://www.fao.org/3/i3531e/i3531e.pdf>
- [28] Allen & Page. (2017). Feeding hens factsheet. Retrieved from <https://www.smallholderfeed.co.uk/wp-content/uploads/2017/09/Feeding-Hens-Factsheet.pdf>
- [29] Organic Feeds. (2023, March 14). How much to feed broiler chickens. Retrieved from <https://organicfeeds.com/how-much-to-feed-broiler-chickens/>
- [30] Aviagen. (2017). Ross 308 broiler manual. Retrieved from <https://kenanaonline.com/files/0070/70697/ross308%20Manual%20broiler.pdf>
- [31] FeedStrategy. (2023). Traditional US broiler feed formulations. Retrieved from <https://www.feedstrategy.com/animal-feed-formulations-library/traditional-us-broiler-feed-formulations>
- [32] CIC Feeds (PVT) LTD. (2023). Chick booster, Broiler starter, Broiler finisher composition.
- [33] National Dairy and Veterinary Sciences Institute. (n.d.). Nutrient requirement and feeding standard for poultry. Retrieved from <https://www.ndvsu.org/images/StudyMaterials/Nutrition/Nutrient-requirement-and-feeding-standard.pdf>

- [34] Mavromichalis, I. (2023). EU red layers development phase. FeedStrategy. Retrieved from <https://www.feedstrategy.com/animal-feed-formulations-library/eu-red-layers-development-phase>
- [35] Mavromichalis, I. (2023). EU red layers production phase. FeedStrategy. Retrieved from <https://www.feedstrategy.com/animal-feed-formulations-library/eu-red-layers-production-phase>
- [36] National Dairy and Veterinary Sciences Institute. (n.d.). Nutrient requirement and feeding standard for poultry. Retrieved from <https://www.ndvsu.org/images/StudyMaterials/Nutrition/Nutrient-requirement-and-feeding-standard.pdf>
- [37] Zulkifli, N. F., Seok-Kian, A. Y., Seng, L. L., Mustafa, S., Kim, Y. S., & Shapawi, R. (2022). Nutritional value of black soldier fly (*Hermetia illucens*) larvae processed by different methods. *PLOS One*, 17(2), e0263924. doi: 10.1371/journal.pone.0263924
- [38] Suryati, T., Julaeha, E., Farabi, K., Ambarsari, H., & Hidayat, A. T. (2023). Lauric acid from the black soldier fly (*Hermetia illucens*) and its potential applications. *Sustainability*, 15(13), 10383. <https://doi.org/10.3390/su151310383>
- [39] Ramos-Bueno, R.P.; Gonzalez-Fernandez, M.J.; Sanchez-Muros-Lozano, M.J.; Barroso, F.G.; Guil-Guerrero, J.L. Fatty acid profiles and cholesterol content of seven insect species assessed by several extraction systems. *Eur. Food Res. Technol.* 2016, 242, 1471–1477.
- [40] Ewald, N.; Vidakovic, A.; Langeland, M.; Kiessling, A.; Sampels, S.; Lalander, C. Fatty acid composition of black soldier fly larvae (*Hermetia illucens*)—Possibilities and limitations for modification through diet. *Waste Manag.* 2020, 102, 40–47.
- [41] Liu, X.; Chen, X.; Wang, H.; Yang, Q.; Rehman, U.K.; Li, W.; Cai, M.; Li, Q.; Mazza, L.; Zhang, J.; et al. Dynamic changes of nutrient composition throughout the entire life cycle of black soldier fly. *PLoS ONE* 2017, 12, e0182601.
- [42] Zheng, L.; Hou, Y.; Li, W.; Yang, S.; Li, Q.; Yu, Z. Biodiesel production from rice straw and restaurant waste employing black soldier fly assisted by microbes. *Energy* 2012, 47, 225–229.
- [43] Leong, S.Y.; Kutty, S.R.; Malakahmad, A.; Tan, C.K. Feasibility study of biodiesel production using lipids of *Hermetia illucens* larva fed with organic waste. *Waste Manag.* 2016, 47, 675–680.
- [44] Suryati, T., Julaeha, E., Farabi, K., Ambarsari, H., & Hidayat, A. T. (2023). Lauric acid from the black soldier fly (*Hermetia illucens*) and its potential applications. *Sustainability*, 15(13), 10383. <https://doi.org/10.3390/su151310383>
- [45] Liu, Z., Li, Y., Ren, H., Zhang, J., Zhao, X., Li, J., Wang, J., Zhang, W., & Wang, X. (2021). The effects of dietary black soldier fly larvae meal on the growth performance, carcass yield, meat quality, and intestinal health of broilers. *Journal of Animal Science*, 99(12), skab375
- [46] Raman, S. S., Stringer, L. C., Bruce, N. C., & Chong, C. S. (2022). Opportunities, challenges and solutions for black soldier fly larvae-based animal feed production. *Journal of Cleaner Production*, 373, 133802.
- [47] Amrul, N. F., Kabir Ahmad, I., Ahmad Basri, N. E., Suja, F., Abdul Jalil, N. A., & Azman, N. A. (2022). A review of organic waste treatment using black soldier fly (*Hermetia illucens*). *Sustainability*, 14(8), 4565. <https://doi.org/10.3390/su14084565>
- [48] Khan, R. A., Naz, S., Rehman, F. U., Farooq, M. Z., Sardar, R., & Nisa, M. (2017). Nutritional and therapeutic potential of black soldier fly (*Hermetia illucens*) larvae: A review. *Journal of Entomology and Zoology Studies*, 5(5), 1801-1807. Retrieved from <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5532560/>
- [49] Khan, R. A., Naz, S., Rehman, F. U., Farooq, M. Z., Sardar, R., & Nisa, M. (2017). Nutritional and therapeutic potential of black soldier fly (*Hermetia illucens*) larvae: A review. *Journal of Entomology and Zoology Studies*, 5(5), 1801-1807. Retrieved from <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5532560/>
- [50] Putra, R., & Safa'at, N. (2020). Study on Sex Determination and Impact of Sex Ratio to Reproduction Success in Black Soldier Fly. *Jurnal Biodjati*, 5(2), 191-198. doi:10.15575/biodjati.v5i2.9472
- [51] Romano, N., Fischer, H., & Egnew, N. (2020). Color and sugar preferences of adult black soldier fly (*Hermetia illucens*) (Diptera: Stratiomyidae) for feeding and oviposition. *Journal of Environmental Biology*, 41(5), 1132-1137. doi:10.22438/jeb/41/5(SI)/MS\_03
- [52] Handayani, D., Naldi, A., Larasati, R.R.N.P., Khaerunnisa, N., & Budiarmaka, D.D. (2021). Management of increasing economic value of organic waste with Maggot cultivation. *IOP Conf. Ser. Earth Environ. Sci.* 2021, 716, 012026.
- [53] EchoCommunity. (2023, October 24). Black soldier fly larvae as animal feed: A guide to production and use. Retrieved from <https://www.echocommunity.org/en/resources/86a8212a-a048-41c6-ab31-536c111e75ac>
- [54] De Souza Vilela, J., Andronicos, N. M., Kolakshyapati, M., Hilliar, M., Sibanda, T. Z., Andrew, N. R., ... & Ruhnke, I. (2022). Black soldier fly larvae in broiler diets improve broiler performance and modulate the immune system. *Animal Nutrition*, 8(1), 1-11. <https://doi.org/10.1016/j.aninu.2022.01.005>